

*Schwind - A. R. S.*

National Aeronautics and Space Administration  
Goddard Space Flight Center  
Contract No. NAS-5-3760

ST — SP — 10 426

NASA TT F-9913

ON SOME CHARACTERISTICS OF RADIO EMISSION SOURCES  
CONNECTED WITH THE ACTIVE REGIONS  
OF THE SUN

by  
A. R. Abbasov  
(USSR)

FACILITY FORM 602

<b>N 66 - 13300</b>	
(ACCESSION NUMBER)	(THRU)
<u>5</u>	<u>1</u>
(PAGES)	(CODE)
	<u>30</u>
(NASA CR OR TMX OR AD NUMBER)	(CATEGORY)

GPO PRICE \$ \_\_\_\_\_

CFSTI PRICE(S) \$ \_\_\_\_\_

Hard copy (HC) 1.00

Microfiche (MF) .50

ff 653 July 65

14 December 1965

ON SOME CHARACTERISTICS OF RADIO EMISSION SOURCES  
CONNECTED WITH THE ACTIVE REGIONS  
OF THE SUN


Vestnik Leningradskogo Universiteta  
No. 14, vyp. 4, 140-142,  
Seriya Matematiki, Mekhaniki i Astronomii  
LENINGRAD, 1965.

by A. R. Abbasov

SUMMARY

This is a preliminary communication on some of the characteristics of radio emission sources connected with the active regions of the Sun. The present observations were conducted during low solar activity. We utilized for their processing the results of observations at IHH and Nagoya stations; they allowed to conclude that a decrease in the S-component of radiation flux is observed prior to intense radiobursts but the flux increases after the bursts, since the sources linked with sunspots are directional.

13300



\*  
\* \*

Radio emission sources above the active regions of the Sun are currently subject to intensive studies. A great number of characteristics of the sources are already known more or less reliably, but some contradictory data still exist in relation to some of them (see [1, 2]). The main issue here is the dependence of source's radio emission flux  $F_S$  on its position on the solar disk when the wavelength  $\lambda < 10$  cm.

Statistical research [3 - 5] made evident the notable weakening of  $F_S$  when the emission source shifts from the center toward the limb of the disk. There exist also observations of separate sources [3, 6, 7], which confirm such a conclusion. On the other hand, certain observations on radiotelescopes with high angular resolution [8 - 10] do not show any notable dependence of  $F_S$  on the distance  $\theta$  from the source to the center of the disk. At the same time the characteristic of the source, referred to above, is of particular importance for resolving the question of its emission mechanism.

\* O NEKOTORYKH KHA-  
RAKTERITIKAKH ISTOCHNIKOV RADIOIZLUCHENIYA SVYAZAN-  
NYKH S AKTIVNYMI OBLASTYAMI NA SOLNTSE.

There is a possibility during the current period of solar activity decrease, of obtaining supplementary information on the character of the dependence  $f(\theta)$  for solitary sources. To that effect we may utilize the routine observations of solar radio emission of [11], brought out in detail in the compilations obtained by the International Center of Data Assembly in Moscow. It is possible to select in the period under consideration those time intervals when there is an active region remaining isolated during the entire period of its shift from  $\theta = 0^\circ$  to  $\theta = 90^\circ$  and for a longer time, that is, not less than two days after that. Taking advantage of the data of optical observations [12], we succeeded in separating for 1962 and 1963 six cases satisfying the imposed requirements (group No. 36 for 1962 and groups No. 43, 46, 101 and 109 for 1963).

The group No. 109 had to be excluded from analysis on account of the great number of radiobursts at  $\theta > 60^\circ$ , which gave a residual increase of  $F_s$ , strongly distorting the real value of the slowly varying emission




Fig. 1




Fig. 2

Radio emission of sources above the remaining five groups was investigated as follows. Using the method of least squares and by way of extrapolation to zero of flocculus and spot areas for the given month, the level  $F_q$  of the "quiet" Sun was determined. Then, the source's emission flux  $F_s = F_f - F_q$  was found by the difference between the daily integral fluxes of Sun's radio emission ( $F_s$ ) and this quiet level. Dividing the flux  $F$  incident upon the spot area ( $S_q$ ) for each day at their shift from the center to the limb, it is possible to find the dependence of source's specific emission on the angular distance  $\theta$  of the spots to the center of the disk (Fig. 1). The results obtained were found to be in agreement with the data of [3 - 7, 15], but departed from the data [8 - 10].

The behavior of sources' emission flux prior to intense radiobursts constituted another characteristic of their radio emission above active regions, of particular interest for the Sun-Earth problem. It is well known that there exists a certain relationship between the characteristics of sources' slowly varying component with bursts, consisting for example in the increase of probability of burst onset either at decrease in the dimensions of the source [13], or in the variation of source's emission spectrum [14].

The dependence of the flux  $F_S$  on  $t$  at  $t < t_b$  ( $t_b$  being the onset time of the burst) was difficult to find at high solar activity, when we had to separate from the aggregate flux of all sources that in whose region the burst was observed. Therefore, the relationship between  $F_S$  and  $t_b$  must be investigated only in four cases (type-C bursts). These cases are plotted in Fig. 2, from which it follows that, prior to the occurrence of the burst there is observed a clearly defined decrease in the emission flux  $F_S$  of the source. Such an effect was characteristic in all cases when the emission flux exceeded the solar flux during the burst by 1000 - 700 %. Prior to the burst the flux decreased by 2.5 - 7 % of the Sun's flux. After the burst the flux remained increased by about 10 to 20 % of the Sun's flux for some 8 - 10 hours.

It is possible that the above-presented method of forecasting intense bursts may also be applied at increased solar activity provided we separate sources are tracked by radiotelescopes with high angular resolution.

The author considers it to be its duty to express his gratitude to A. P. Molchanov and A. S. Grebinskiy for their valuable advice when discussing the results.

\*\*\*\*\* THE END \*\*\*\*\*

Contract No. NAS-5-3760  
Consultants & Designers, Inc.  
Arlington, Virginia

Translated by ANDRE L. BRICHANT  
on 13 December 1965

#### REFERENCES

1. - KURS ASTROFIZIKI I ZVEZDNOY ASTRONOMII. - Gl. IX. - Pod red. A. A. Mikhaylova. M., NAUKA, 1964.
2. - V. V. ZHELEZNIKOV. - Radioizlucheniye solntsa i planet (Sun's and Planets' Radioem.) M., NAUKA, 1965.
3. - A. P. MOLCHANOV, R. D. DAGKESAMANSKIY. - Soln. dannyye (Solar Data), 5, 77, 1960
4. - KIN-AKI KAWABATA. - Publ. Astr. Soc. Japan, 12, No. 4, 513, 1960.
5. - B. VOKUA. - Radioastronomiya (Parizhskiy simp. 1958) 143 M. IL, 1961.
6. - M. GUTMAN, ZH. L. STEYNBERG. - Ib. 123, M., 1961.
7. - W. N. CHRISTIANSEN, A. BOISCHOT., a.u. Ann. d'Astrophysique 23, No. 1, 75, 1960.
8. - V. V. VITKEVICH, A. D. KUZ'MIN, A. E. SALOMONOVICH et al, Radioastronom. Parizhskiy simp. 1958, 129, M., IL 1961.
9. - A. P. MOLCHANOV, N. L. KAYDANOVSKIY, N. G. PETROVA. - Sol. d. 9, 67, 1964.
10. - V. Ya. GOL'NEV, V. N. ISKHANOVA, et al. - Ib. No. 9, 67, 1964.
11. - Quartely Bull. Solar Data. - Zurich.
12. - Byull. Sol. dannyye No. 3, 1962; No. 9, 10, 1963.
13. - M. R. KUNDYU. - Radioastronomiya (Parizhsk. simpos. 1958) 219., M., IL 1961.
14. - H. TANAKA. Proc. res. U. Inst. atmosph. Nagoya, 11, 41, 1964.
15. - U. V. KHAYGIL'DIN. - Astronom. Zh., 26, 2, 1964.

D I S T R I B U T I O NG O D D A R D S. F. C.

100 CLARK, TOWNSEND  
 110 STROUD  
 400 BOURDEAU  
 110 MEREDITH  
 611 MC DONALD  
     ABRAHAM  
     BOLDT  
     WILLIAMS  
 612 HEPPNER  
     NESS  
 613 KUPPERIAN (3)  
 614 WHITE  
     HALLAM  
     BEHRING  
     WOLFF  
     FROST  
 615 BAUER  
     GOLDBERG  
     STONE  
 640 HESS (3)  
     MEAD  
     CAMERON  
 630 GI for SS (5)  
 252 LIBRARY  
 256 FREAS

N A S A H Q S.

SS NEWELL, NAUGLE  
 SG MITCHELL  
     ROMAN  
     SMITH  
     DUBIN  
 SL LIDDEL  
     FELLOWS  
     BRUNK  
     HIPSHER  
     HOROWITZ  
 SM FOSTER  
     ALLENBY  
     GILL  
     BADGLEY  
 RR KURZWEG  
 RTR  
     NEILL  
 ATSS  
     SCHWIND  
     ROBBINS  
 WX SWEET

O T H E R C E N T E R SAMES

LIBRARY (3)  
 SONETT (5)

JPL

SNYDER  
 NEWBURN  
 WYCKOFF

LANGLEY

160 ADAMSON  
 213 KATZOFF  
 235 SEATON  
 185 WEATHERWAX

UCLA

COLEMAN

U. C. at BERKELEY

WILCOX

M. I. T.

BARRETT

U. MICHIGAN

ARNOLD  
 HADDOCK

S. W. C. A. S.

JOHNSON